AMENDMENT

In the claims

Please cancel claims 4, 5, and 7-11. Please replace claims 6 and 12-14 with the following claims:

A method of making an oligonucleotide, the method comprising:

providing a template oligonucleotide comprising a sequence of nucleotides, the template comprising at least one non-standard nucleotide at a preselected site in the sequence;

contacting the template with a mixture of nucleotide triphosphates, the mixture comprising nucleotide triphosphates that are complementary to the nucleotides of the template, wherein the nucleotide triphosphate complementary to the non-standard nucleotide at the preselected site comprises a derivatized nucleotide; and

forming an oligonucleotide complementary to a portion of the template by enzymatic polymerization of the nucleotide triphosphates in a sequence complementary to the portion of the template, wherein the non-standard nucleotide at the preselected site is iso-G or iso-C.

A method of making an oligonucleotide, the method comprising:

providing a template oligonucleotide comprising a sequence of nucleotides, the template comprising at least one non-standard nucleotide at a preselected site in the sequence;

contacting the template with a mixture of nucleotide triphosphates, the mixture comprising nucleotide triphosphates that are complementary to the nucleotides of the template, wherein the nucleotide triphosphate complementary to the non-standard nucleotide at the preselected site comprises a derivatized nucleotide; and

forming an oligonucleotide complementary to a portion of the template by enzymatic polymerization of the nucleotide triphosphates in a sequence complementary to the portion of the template, wherein the enzymatic polymerization

F

۴

F is performed with
A comprises a DNA polymerase selected from the group consisting of AMV reverse transcriptase, 74-DNA polymerase, and Klenow fragment of DNA polymerase 1.

The method of claim 1/2, wherein the DNA polymerase comprises Klenow fragment of DNA polymerase I.

14. A method of making an oligonucleotide, the method comprising:

providing a template oligonucleotide comprising a sequence of nucleotides, the template comprising at least one non-standard nucleotide at a preselected site in the sequence;

contacting the template with a mixture of nucleotide triphosphates, the mixture comprising nucleotide triphosphates that are complementary to the nucleotides of the template, wherein the nucleotide triphosphate complementary to the non-standard nucleotide at the preselected site comprises a derivatized nucleotide comprising a radiolabel; and

forming an oligonucleotide complementary to a portion of the template by enzymatic polymerization of the nucleotide triphosphates in a sequence complementary to the portion of the template, wherein the enzymatic polymerization is performed with a comprises T7 RNA polymerase.

-3-